

AD _____

Award Number: DAMD17-01-1-0132

TITLE: To Ascertain Distinctive Gene Expression Patterns for the
Prediction of Docetaxel (Taxotere) Chemosensitivity or
Chemoresistance in Human Breast Cancer

PRINCIPAL INVESTIGATOR: Jenny C. Chang, M.D.

CONTRACTING ORGANIZATION: Baylor College of Medicine
Houston, Texas 77030

REPORT DATE: October 2002

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are
those of the author(s) and should not be construed as an official
Department of the Army position, policy or decision unless so
designated by other documentation.

20030319 042

REPORT DOCUMENTATION PAGEForm Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE October 2002	3. REPORT TYPE AND DATES COVERED Annual (17 Sep 01 - 16 Sep 02)	
4. TITLE AND SUBTITLE To Ascertain Distinctive Gene Expression Patterns for the Prediction of Docetaxel (Taxotere) Chemosensitivity or Chemoresistance in Human Breast Cancer			5. FUNDING NUMBERS DAMD17-01-1-0132	
6. AUTHOR(S) Jenny C. Chang, M.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Baylor College of Medicine Houston, Texas 77030 E-Mail: jcchang@breastcenter.tmc.edu			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited				12b. DISTRIBUTION CODE
13. ABSTRACT (Maximum 200 Words) Chemotherapy is of proven benefit in reducing the risk of death for a subset of patients with early breast cancer, but doctors have problems deciding exactly who should receive this therapy, and which therapy will be most effective for a given patient. As a result, some patients needlessly receive chemotherapy. Even in those patients who clearly require chemotherapy, doctors cannot identify those patients whose tumors might not be responsive to a particular chemotherapy drug. Chemotherapy is also associated with high costs and toxicity including nausea, vomiting, damage to nerves, etc. and increased risk of infections that are sometimes life threatening. The emerging cDNA array technology provides a means to comprehensively appreciate genetic variations in different breast tumors, and may be utilized as a test for chemotherapy sensitivity. Taxotere has one of the highest response rates in breast cancer, and is widely prescribed for the treatment of breast cancer. The aims of this study are therefore, to investigate and validate differential gene expression patterns from core biopsies from patients whose breast tumors either shrank after Taxotere chemotherapy, or failed to respond. These genes whose expression patterns are associated with Taxotere response could be used to create a simple test of predictor genes to help doctors treat breast cancer more effectively.				
14. SUBJECT TERMS breast cancer				15. NUMBER OF PAGES 9
				16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

Table of Contents

Cover.....	1
SF 298.....	2
Introduction.....	4
Body.....	5
Key Research Accomplishments.....	6
Reportable Outcomes.....	7
Conclusions.....	7
References.....	7
Appendices.....	

Commander
U.S. Army Medical Research and Materiel Command
504 Scott Street
Fort Detrick, MD 21702-5012

Subject: Annual Report for Award Number DAMD17-01-1-0132

Introduction

Optimal systemic treatment after breast cancer is the most crucial factor in reducing mortality in women with breast cancer. Adjuvant chemotherapy and hormonal treatment both reduce the risk of death in breast cancer patients. However, while estrogen receptors status predicts for response to hormone treatments, there are no clinically useful predictive markers for chemotherapy response. All eligible women are therefore treated in the same manner, even though de novo drug resistance will result in treatment failures in many breast cancer patients. Currently, there are no methods available to distinguish those patients who are likely to respond to specific chemotherapies, and given the accepted practice of prescribing adjuvant treatment to most patients, even if the average expected benefit is slow, the selection of appropriate patients represents a major advance in the clinical management of breast cancer today.

We therefore set out to identify gene expression patterns in primary breast cancer specimens that might predict response to Taxotere. Neoadjuvant chemotherapy allows for the sampling of the primary tumor for gene expression analysis and for direct assessment of response to chemotherapy by following changes in tumor size during the first few months of treatment. Hence, neoadjuvant chemotherapy provides an ideal platform to rapidly discover predictive markers of chemotherapy response.

In this present study, we hypothesize that high throughput quantitation of gene expression, it is possible to assess thousands of genes simultaneously, and that these expression patterns in different breast cancers might correlate with and thereby predict response to treatment. The purpose of this study was to (1) demonstrate that sufficient RNA could be obtained from core biopsies to access gene expression, (2) to identify groups of genes that could be used to distinguish primary breast cancers to responsive or resistance to Ttaxotere, and (3) to identify gene pathways that could be important in a mechanism of action of taxotere.

Body of Research

From September 17, 2001 to September 16, 2002, we have recruited 42 patients with locally advanced breast cancer. Core biopsies were obtained from the primary breast cancers before commencement of neoadjuvant chemotherapy. Clinical responses before and after four cycles of chemotherapy were measured in all primary breast cancers.

A total of 6 core biopsies were obtained from each primary cancer. Two core biopsy specimens were transferred immediately to liquid nitrogen and snap frozen at -80°C . Each core biopsy measured approximately 1 cm x 1 mm. As these core biopsies were too small for micro dissection, we ascertained the tumor cellularity of the pretreatment core biopsies. In general, the core biopsies showed good tumor cellularity with median tumor cellularity of 75% (range 40-100%). Each core biopsy yielded 3-6 mg of total RNA, which is more than sufficient to generate approximately 20 mg of label cRNA needed for hybridization with the Affymetrix U95Av2 Genechip, using the manufacturer's standard protocols. To date, we have analyzed 24 out of the 42 collected core biopsy specimens. We have begun initial exploratory analysis. We compared

the expression data in sensitive and resistant tumors to identify gene significance differentially expressed between the two groups. We applied filtering to eliminate genes with uniformly low expression or whose expression did not vary significantly across the samples retaining approximately 1,600 genes. We then applied t-test after log transformation, to select discriminatory genes. To date, we have selected 92, 300, 551 genes as differentially expresses at p values of 0.001, 0.01, 0.05, respectively.

In the 92 gene list, among the genes overexpressed in the resistant cluster, most are involved in transcriptional regulation, signal transduction, or have unknown functions. In the sensitive tumors, some are involved in signal transduction and cell cycle, cytoskeleton and adhesion processes, protein transport, protein modification, stress and apoptosis or have unknown functions. We recently begun to confirm the expression measurements obtained from the affymetrics chips with values from semiquantitative RT-PCR. We have done 15 genes and compared their measurements with QRT PCR. Significantly correlation was seen between the two methods.

Key Research Accomplishments

Three abstracts have been submitted and accepted for international meetings. Two were submitted to the San Antonio Breast Cancer Symposium in 2001 and 2002. This abstract was also submitted to the ASCO meeting in Florida in 2002. This was selected for oral presentation. A manuscript is underway to discuss the preliminary findings of this study. This study was also selected for a plenary presentation in the Era of Hope Meeting in Florida in 2002.

Reportable Outcomes

1. Genetic markers for response to neoadjuvant therapy: Array based gene expression profiling from serial biopsies. EC Wooten, **J Chang**, SG Hilsenbeck. 24th Annual San Antonio Breast Cancer Symposium, San Antonio, Texas (abstract 236), December 2001.
2. Gene expression profiles from breast cancer core biopsies predict therapy to response. EC Wooten, **J Chang**, SG Hilsenbeck. *Proceedings of the American Association for Cancer Research* 43, abstract 450, March 2002.
3. Gene expression profiles for doxytaxcil chemosensitivity. **J Chang**, EC Wooten and R Elledge. ASCO 28th Annual Meeting, abstract 1700, May 2002.

Conclusions

We have determined that 1) sufficient RNA can be obtained from core needle biopsies to hybridize Affymetrix GeneChips for assessment of gene expression patterns 2) differential gene expression patterns exist that can distinguish resistant *versus* sensitive tumors. We will continue our current experiments to further increase patient recruitment to define and refine patterns of resistance and sensitivity.

References

1. EBCTG. Systemic treatment of early breast cancer by hormonal, cytotoxic or immune therapy: 133 randomised trials involving 31,000 recurrences and 24,000 deaths among 75,000 women. *Lancet* **339**, 1-15, 71-85 (1992).
2. The Early Breast Cancer Trialists' Collaborative Group, E. Tamoxifen for early breast cancer: an overview of the randomised trials. *Lancet* **351**, 1451-1467 (1998).
3. EBCTG. Polychemotherapy for early breast cancer: an overview of the randomised trials. *Lancet* **352**, 930-942 (1998).

4. Hortobagyi, G.N. Docetaxel in breast cancer and a rationale for combination therapy. *Oncology* **11**, 11-15 (1997).
5. Chan, S. et al. Prospective randomized trial of docetaxel versus doxorubicin in patients with metastatic breast cancer. The 303 Study Group. *J Clin Oncol* **17**, 2341-54. (1999).
6. Nabholz, J.M. et al. A phase III trial comparing docetaxel (T), doxorubicin (A) and cyclophosphamide (C) (TAC) to (FAC) as first line chemotherapy for patients with metastatic breast cancer. *Proceedings of the American Society of Clinical Oncologists* **20**, 22a (2001).
7. Henderson IC, B.D., Demetri G, et al. Improved disease free survival and overall survival from the addition of sequential paclitaxel but not from escalation of doxorubicin in the adjuvant chemotherapy of patients with node-positive primary breast cancer. *Proc Am Soc Clin Onco* **17**, 101 (1998).
8. Aapro, M.S. Adjuvant therapy of primary breast cancer: a review of key findings from the 8th international conference, St. Gallen. *The Oncologist* **6**, 376-385 (2001).
9. Fisher, B. et al. Effect of preoperative chemotherapy on the outcome of women with operable breast cancer. *Journal of Clinical Oncology* **16**, 2672-2685 (1998).
10. Mamounas, E.P. Preoperative doxorubicin plus cyclophosphamide followed by preoperative or postoperative docetaxel. *Oncology* **11**, 37-40 (1997).
11. Schena, M., Shalon, D., Davis, R.W. & Brown, P.O. Quantitative monitoring of gene expression patterns with a complementary DNA microarray. *Science* **270**, 467-470 (1995).
12. Khan, J. et al. Gene expression profiling of alveolar rhabdomyosarcoma with cDNA microarrays. *Cancer Research* **58**, 5009-5013 (1998).
13. Lockhart, D.J., Dong, H., Byrne, M.C. & et al. Expression monitoring by hybridization to high-density oligonucleotide arrays. *Nature Biotechnology* **14**, 1675-1680 (1996).
14. Sgroi, D.C. et al. In vivo gene expression profile analysis of human breast cancer progression. *Cancer Research* **59**, 5656-5661 (1999).
15. Perou, C.M. et al. Distinctive gene expression patterns in human mammary epithelial cells and breast cancers. *Proceedings of the National Academy of Sciences of the United States of America* **96**, 9212-9217 (1999).
16. Li, C. & Wong, W.H. Model-based analysis of oligonucleotide arrays: expression index computation and outlier detection. *Proceedings of the National Academy of Sciences of the United States of America* **98**, 31-36 (2001).
17. Schadt, E.E., Li, C. & Wong, W.H. Analyzing high-density oligonucleotide gene expression array data. *Journal of Cellular Biochemistry* **80**, 192-202 (2001).
18. Tusher, V.G., Tibshirani, R. & Chu, G. Significance analysis of microarrays applied to the ionizing radiation response. *Proc Natl Acad Sci U S A* **98**, 5116-21. (2001).
19. Hakak, Y. et al. Genome-wide expression analysis reveals dysregulation of myelination-related genes in chronic schizophrenia. *Proceedings of the National Academy of Sciences of the United States of America* **98**, 4746-4751 (2001).
20. Bowcock, A.M. et al. Insights into psoriasis and other inflammatory diseases from large-scale gene expression studies. *Human Molecular Genetics* **10**, 1793-1805 (2001).
21. Van Poznak, et al. Assessment of molecular markers of clinical sensitivity to single agent taxane therapy for metastatic breast cancer. *Journal of Clinical Oncology* **20(9)**, 2319-2326 (2002).

22. Hansen, R.K. et al. Hsp27 overexpression inhibits doxorubicin-induced apoptosis in human breast cancer cells. *Breast Cancer Res Treat* **56**, 187-96. (1999).